

Air Pollution and Mortality in Seven Million Adults: The Dutch Environmental Longitudinal Study (DUELS)

Paul H. Fischer,¹ Marten Marra,¹ Caroline B. Ameling,¹ Gerard Hoek,² Rob Beelen,^{1,2} Kees de Hoogh,^{3,4,5} Oscar Breugelmans,¹ Hanneke Kruize,¹ Nicole A.H. Janssen,¹ and Danny Houthuijs¹

¹National Institute for Public Health and the Environment, Bilthoven, the Netherlands; ²Institute for Risk Assessment Sciences, Utrecht University, Utrecht, the Netherlands; ³MRC-PHE Centre for Environment and Health, Department of Epidemiology and Biostatistics, Imperial College London, London, United Kingdom; ⁴Swiss Tropical and Public Health Institute, Basel, Switzerland; ⁵University of Basel, Basel, Switzerland

BACKGROUND: Long-term exposure to air pollution has been associated with mortality in urban cohort studies. Few studies have investigated this association in large-scale population registries, including non-urban populations.

OBJECTIVES: The aim of the study was to evaluate the associations between long-term exposure to air pollution and nonaccidental and cause-specific mortality in the Netherlands based on existing national databases.

METHODS: We used existing Dutch national databases on mortality, individual characteristics, residence history, neighborhood characteristics, and national air pollution maps based on land use regression (LUR) techniques for particulates with an aerodynamic diameter $\leq 10 \mu\text{m}$ (PM_{10}) and nitrogen dioxide (NO_2). Using these databases, we established a cohort of 7.1 million individuals ≥ 30 years of age. We followed the cohort for 7 years (2004–2011). We applied Cox proportional hazard models adjusting for potential individual and area-specific confounders.

RESULTS: After adjustment for individual and area-specific confounders, for each $10\text{-}\mu\text{g}/\text{m}^3$ increase, PM_{10} and NO_2 were associated with nonaccidental mortality [hazard ratio (HR) = 1.08; 95% CI: 1.07, 1.09 and HR = 1.03; 95% CI: 1.02, 1.03, respectively], respiratory mortality (HR = 1.13; 95% CI: 1.10, 1.17 and HR = 1.02; 95% CI: 1.01, 1.03, respectively), and lung cancer mortality (HR = 1.26; 95% CI: 1.21, 1.30 and HR = 1.10 95% CI: 1.09, 1.11, respectively). Furthermore, PM_{10} was associated with circulatory disease mortality (HR = 1.06; 95% CI: 1.04, 1.08), but NO_2 was not (HR = 1.00; 95% CI: 0.99, 1.01). PM_{10} associations were robust to adjustment for NO_2 ; NO_2 associations remained for nonaccidental mortality and lung cancer mortality after adjustment for PM_{10} .

CONCLUSIONS: Long-term exposure to PM_{10} and NO_2 was associated with nonaccidental and cause-specific mortality in the Dutch population of ≥ 30 years of age.

CITATION: Fischer PH, Marra M, Ameling CB, Hoek G, Beelen R, de Hoogh K, Breugelmans O, Kruize H, Janssen NA, Houthuijs D. 2015. Air pollution and mortality in seven million adults: the Dutch Environmental Longitudinal Study (DUELS). *Environ Health Perspect* 123:697–704; <http://dx.doi.org/10.1289/ehp.1408254>

Introduction

Long-term exposure to air pollution has been associated with mortality in several cohort studies (Abbey et al. 1999; Beelen et al. 2014; Brunekreef et al. 2009; Carey et al. 2013; Cesaroni et al. 2013; Chen et al. 2013; Crouse et al. 2012; Dockery et al. 1993; Hales et al. 2012; Huss et al. 2010; Pope et al. 1995; Yap et al. 2012; Zeger et al. 2008). Although the evidence is increasing, heterogeneity in size of effect estimates between cohort studies has been identified (Hoek et al. 2013).

Cohort studies specifically designed for investigating individual risk factors are time consuming, labor intensive, often limited in size, and relatively costly. To overcome these disadvantages, recent studies have linked existing national databases of air pollution, nonaccidental mortality, individual characteristics, and residential history to assess the relationships between air pollution and mortality more efficiently (Carey et al. 2013; Cesaroni et al. 2013; Chen et al. 2013; Crouse et al. 2012; Hales et al. 2012; Huss

et al. 2010; Zeger et al. 2008). The aim of our study was to use existing national databases to evaluate the associations of long term exposure to air pollution [particulates with an aerodynamic diameter $\leq 10 \mu\text{m}$ (PM_{10}) and nitrogen dioxide (NO_2)] with nonaccidental and cause-specific mortality in a cohort of 7.1 million Dutch residents.

Methods

The study cohort. In the Netherlands, population statistics are compiled by Statistics Netherlands (<http://www.cbs.nl/en-GB/menu/home/default.htm>) and are based on digital municipal population registers (Prins 2000). This registration system is known as the GBA (Gemeentelijke Basis Administratie), the municipal basic registration of population data. The GBA was implemented on 1 October 1994.

Statistics Netherlands combines the data from the GBA into a longitudinal file for each individual registered in the GBA (de Bruin et al. 2004). These records start on 1 January 1995. Changes in demographic

attributes (e.g., death, address, marital status, emigration) are updated yearly by adding additional information on the nature and the date of the change. In these files, the individual identification number of the GBA is replaced by a meaningless, but unique, identification number. This identification number is used to enrich the individual files with information from other central data sources maintained by Statistics Netherlands such as the social statistical database, which contains, among others, data from the tax authorities and about employment status (Arts and Hoogteijling 2002).

From the database with the longitudinal files of all Dutch inhabitants, we selected all individuals of ≥ 30 years of age on 1 January 2004, living at the same residential address since 1 January 1999. We used data about sex, age, marital status, and region of origin. The data about origin distinguishes between

Address correspondence to P.H. Fischer, Department for Environmental Health, Centre for Sustainability, Environment and Health, National Institute of Public Health and the Environment, Bilthoven, the Netherlands. Telephone: 31 30 2743315. E-mail: paul.fischer@rivm.nl

Supplemental Material is available online (<http://dx.doi.org/10.1289/ehp.1408254>).

The research leading to these results was performed in close cooperation with Statistics Netherlands, which provided the national registry data and facilitated the statistical analyses. F. Janssen (Population Research Centre, Faculty of Spatial Sciences, University of Groningen) kindly provided the data on regional age-standardized smoking attributable mortality fractions; D. de Zwart (RIVM) kindly supported with editing the figures, and W. Swart (RIVM) with the analyses on geo-information systems.

The study was funded by the RIVM Strategic Research Program and The Ministry of Infrastructure and the Environment. The exposure assessment work to create exposure maps for 2001 was supported by grant RGI-137 from the Dutch program Ruimte voor Geoinformatie supported by the Ministry of Housing, Spatial Planning and the Environment. For the sensitivity analyses, we used data collected by Community Health Services (GGD) of GGD Fryslan, GGD Gooi & Vechtstreek, GGD Hollands Midden, GGD Rotterdam-Rijnmond, GGD Zuid-Holland West, GGD Hart voor Brabant, GGD West Brabant, GGD Brabant Zuidoost, GGD Limburg Noord, and GGD Zuid Limburg.

The authors declare they have no actual or potential competing financial interests.

Received: 10 February 2014; Accepted: 6 March 2015; Advance Publication: 11 March 2015; Final Publication: 1 July 2015.

Dutch, Western origin, and non-Western origin. Individuals of non-Western origin are those born in or with a parent born in Africa, Asia (except Japan and Indonesia, who are categorized as “Western origin”), or Latin America. Given the relatively large size of groups within non-Western origin, a distinction is made among those with Turkey, Morocco, and Suriname origin. Furthermore, we enriched the database with standardized disposable household income. This individual socioeconomic indicator is adjusted for differences in household size and composition.

We also used a socioeconomic indicator at four-digit postal code level. These postal code areas comprise on average about 4,000 inhabitants. This social status indicator is derived every 4 years by the Netherlands Institute for Social Research (<http://www.scp.nl/english/>) (Knol 1998). Each postal code area receives a unique ranking for social status according to the income level, unemployment rate, and education level of its inhabitants. The ranking is transformed to a 0–1 scale, with 1 being the lowest possible ranking on social status within the Netherlands. We took the indicator from 2002 and linked it to the cohort through the postal code of the residential addresses.

The follow-up period of the cohort was from 1 January 2004 to 1 January 2011. Subjects were lost to follow-up if their final record in the longitudinal file ended before 1 January 2011 and death was not registered as a reason for termination. Emigration was the main cause of censoring.

The Dutch population registers are intended primarily for municipal administrative purposes. However, many national and nongovernmental organizations benefit from them as well. Given the confidential character of the data, there is no free access to the population registers. Each organization interested in receiving data on a regular basis is given the opportunity to use the data upon request to the Ministry for the Interior. The Ministry decides to which data the organization gets access.

All our analyses were performed within strict privacy rules; that is, only researchers who received a signed permit were allowed to do analyses within a secured environment at our Institute. Before publication, Statistics Netherlands made sure that none of the analysis results showed potential reducibility to the individual level.

Mortality outcomes. A database with mortality data was available from Statistics Netherlands (Harteloh et al. 2010). We selected nonaccidental mortality [International Classification of Diseases, 10th Revision (ICD-10) codes A00–R99], circulatory disease mortality (ICD-10 codes I00–I99), respiratory disease mortality (ICD-10 codes J00–J99), and lung cancer mortality (ICD-10 codes C33–C34). A study of cause-of-death coding

showed high reliability for these specific causes (> 90% for major causes of death such as cancers and acute myocardial infarction and about 85% for respiratory disease mortality) (Harteloh et al. 2010).

Air pollution exposure assessment. We made use of previously published land use regression (LUR) models to produce high-resolution air pollution maps (100 m × 100 m grids) of annual mean concentrations of PM₁₀ and NO₂ in 2001. Details of development and validation of the LUR models are presented elsewhere (Vienneau et al. 2010). Briefly, for both pollutants regression models were derived from annual mean concentrations for the year 2001 based on routine measurement data from the Dutch National Air Quality Monitoring Network (van Elzakker and Buijsman 2014). Predictor variables used for the modeling were traffic, land use, and topography integrated in a geographical information system. Addresses at baseline were linked to the estimated PM₁₀ and NO₂ concentration in the corresponding grid.

We did not assign PM_{2.5} (PM with diameter ≤ 2.5 μm) concentrations to cohort addresses, because in 2001 PM_{2.5} was not measured in the national monitoring network. Two monitoring studies (Cyrys et al. 2003; Eeftens et al. 2012) showed that the spatial variation of PM₁₀ in the Netherlands is largely driven by PM_{2.5} ($R^2 = 0.76$ and 0.72) with a median ratio between PM_{2.5} and PM₁₀ of 0.66 that was stable over time from 2000 to 2009.

Statistical analyses. Statistical analyses were performed with SAS version 9.1 (SAS Institute Inc., Cary, NC, USA). We applied age-stratified Cox proportional hazards regression models to estimate the associations [hazard ratio (HR) and 95% confidence interval (CI)] between (cause-specific) mortality and long-term exposure to PM₁₀ or NO₂. We used 1-year age strata. We analyzed the data with *a*) models adjusting for age and sex (“unadjusted” model); *b*) models adjusted for age, sex, marital status, region of origin, and household income (individual confounder model); *c*) the individual confounder models extended with a socioeconomic status indicator of postal code areas (full model). In addition, we *d*) extended the full model with the second pollutant to analyze the robustness of the one-pollutant estimate when adjusted for the second pollutant. Statistical significance was defined as p -values < 0.05.

We explored nonlinearity in the relationships between PM₁₀ and NO₂ exposure and mortality with natural splines (2 degrees of freedom). We used the likelihood ratio test (LRT) ($p < 0.05$) to compare spline models with linear models. We analyzed these models with R version 2.15.1 (R Core Team 2014).

To assess the sensitivity of our relative risk estimates to missing individual lifestyle factor data, we assessed the association between our air pollution exposure estimates and lifestyle factors in a separate survey of adults across the Netherlands. We obtained data from health surveys from Community Health Services (GGD GHOR Nederland 2014) conducted in 2003–2005. We included data from 11 Community Health Services with available information on self-reported four-digit postal code, age, sex, marital status, level of education, region of origin, smoking, body mass index (BMI), alcohol consumption, and exercise. Criteria for alcohol consumption were defined by a national working group of experts for the purpose of the Community Health Services health surveys. We calculated the age- and sex-adjusted mean PM₁₀ and NO₂ concentrations at four-digit postal code level for different categories of smoking (current smoker, former smoker, never smoker), BMI (< 18.5, 18.5–25, 25–30, > 30), alcohol consumption (different categories of compliance to three criteria for responsible alcohol use), and exercise (compliance to 30 min of moderate exercise per day on at least 5 days per week). Subsequently, we additionally adjusted for the individual

Table 1. Characteristics of the cohort ($n = 7,218,363$) at baseline (2004).

Characteristic/category	n (%)
Sex	
Male	3,444,166 (47.7)
Age (years)	
30–40	1,053,506 (14.6)
40–50	1,765,274 (24.5)
50–60	1,804,214 (25.0)
60–70	1,256,075 (17.4)
70–80	877,444 (12.2)
≥ 80	461,850 (6.4)
Marital status	
Married	4,977,475 (69.0)
Single	924,165 (12.8)
Widowed	639,451 (8.9)
Divorced	528,753 (7.3)
Other	148,519 (2.1)
Origin ^a	
Morocco	67,444 (0.9)
Turkey	84,773 (1.2)
Suriname	91,588 (1.3)
Non-Western	119,538 (1.7)
Western	638,103 (8.8)
Dutch	6,216,917 (86.1)
Mortality (ICD-10 code)	
Total nonaccidental (A00–R99)	668,206 (9.3)
Diseases of the circulatory system (I00–I99)	209,940 (2.9)
Diseases of the respiratory system (J00–J99)	65,132 (0.9)
Lung cancer (C33–C34)	53,735 (0.7)

^aIndividuals of non-Western origin are those born in or with a parent born in Africa, Asia (except Japan and Indonesia, categorized as “Western origin”), or Latin America. Given the relative large size of groups within non-Western origin a distinction is made between Turkey, Morocco, and Suriname origin.

and neighborhood confounders that were also included in the Cox proportional hazard regression models [i.e., marital status, region of origin, individual socioeconomic status (using level of education in place of standardized household income, which was not available), and social status]. In addition to these regression analyses, we calculated the prevalence of the different variables under study for different categories (deciles) of PM₁₀ and NO₂ exposure.

Furthermore, in the full population we additionally adjusted for area-level smoking-related mortality estimated based upon observed lung cancer rates (Janssen and Spiensma 2012).

In addition we assessed effect modification by stratifying our analyses by sex, age (30–65 or > 65 years), socioeconomic status (five categories), and degree of urbanization (five categories). Results are graphically presented.

Results

On 1 January 2004 the total population of the Netherlands was 16,260,465, of whom 9,936,994 had not moved residence in the previous 5 years (61%). Of these, 7,218,363 were of ≥ 30 years of age (73%) and entered the cohort. Table 1 shows the characteristics of the cohort members on 1 January 2004. During the follow-up period until 1 January 2011, 668,206 (9.3%) cohort members died from natural causes. Of these, 209,940 (31.4%) died from diseases of the circulatory system, 65,132 (9.7%) died from diseases of the respiratory system, and 53,735 (8.0%) died from lung cancer (Table 1).

Figure 1 shows maps of the distributions of the estimated PM₁₀ and NO₂ concentrations in the Netherlands for 2001. For the addresses of the cohort members the median PM₁₀ concentration was 29 μg/m³ [5th–95th percentile, 24 μg/m³–32 μg/m³; interquartile range (IQR) = 2.4]; the median NO₂ concentration was 31 μg/m³ (5th–95th percentile, 19 μg/m³–44 μg/m³; IQR = 10.0 μg/m³). We estimated HRs per 10-μg/m³ increase in the pollutant concentration. When expressed per IQR, the estimates for PM₁₀ become smaller because the IQR for PM₁₀ is smaller than for

NO₂. The range (and IQR) in NO₂ concentrations is larger than the range in PM₁₀ concentrations, because NO₂ is more influenced by local (traffic) emissions than PM₁₀, which is more affected by long-range transport.

The correlation (*R*) between PM₁₀ and NO₂ was 0.58.

The results of the Cox proportional hazard analyses are presented in Tables 2 (PM₁₀) and 3 (NO₂). The highest HRs were found for both pollutants and for each category of cause of death for the “unadjusted” model. Adding individual confounders and area level for

Table 2. Hazard ratios (95% CIs) per 10-μg/m³ increase in PM₁₀ levels at the home address in 2001 for nonaccidental mortality, mortality from circulatory diseases, mortality from respiratory diseases, and lung cancer mortality.

PM ₁₀	Nonaccidental	Circulatory diseases	Respiratory diseases	Lung cancer
Unadjusted model	1.13 (1.12, 1.14)	1.11 (1.09, 1.13)	1.18 (1.14, 1.22)	1.30 (1.25, 1.35)
Individual confounders model	1.10 (1.09, 1.11)	1.08 (1.06, 1.10)	1.16 (1.12, 1.20)	1.30 (1.25, 1.35)
Full model	1.08 (1.07, 1.09)	1.06 (1.04, 1.08)	1.13 (1.10, 1.17)	1.26 (1.21, 1.30)
Two-pollutant model	1.04 (1.03, 1.06)	1.09 (1.07, 1.12)	1.16 (1.11, 1.20)	1.09 (1.04, 1.14)

Unadjusted model: adjusted for age and sex. Individual confounders model: adjusted for age, sex, marital status, region of origin, and standardized household income. Full model: adjusted for neighborhood (postal digit) social status, in addition to the individual confounders. Two-pollutant model: adjusted for estimated NO₂/PM₁₀ at the home address in 2001, in addition to the full model covariates.

Table 3. Hazard ratios (95% CIs) per 10-μg/m³ increase in NO₂ levels at the home address in 2001 for nonaccidental mortality, mortality from circulatory diseases, mortality from respiratory diseases, and lung cancer mortality.

NO ₂	Nonaccidental	Circulatory diseases	Respiratory diseases	Lung cancer
Unadjusted model	1.06 (1.06, 1.06)	1.03 (1.03, 1.04)	1.05 (1.04, 1.06)	1.14 (1.13, 1.15)
Individual confounders model	1.04 (1.04, 1.04)	1.01 (1.01, 1.02)	1.03 (1.02, 1.04)	1.12 (1.11, 1.14)
Full model	1.03 (1.02, 1.03)	1.00 (0.99, 1.01)	1.02 (1.01, 1.03)	1.10 (1.09, 1.11)
Two-pollutant model	1.02 (1.02, 1.02)	0.98 (0.98, 0.99)	0.99 (0.98, 1.00)	1.08 (1.07, 1.10)

Unadjusted model: adjusted for age and sex. Individual confounders model: adjusted for age, sex, marital status, region of origin, and standardized household income. Full model: adjusted for neighborhood (postal digit) social status, in addition to the individual confounders. Two-pollutant model: adjusted for estimated NO₂/PM₁₀ at the home address in 2001, in addition to the full model covariates.




Figure 1. Distributions of the estimated PM₁₀ and NO₂ concentrations in the Netherlands for the year 2001 (modeled with land use regression models).

socioeconomic status reduced the magnitude of the associations.

We estimated the following associations for a 10- $\mu\text{g}/\text{m}^3$ increase in exposure to PM_{10} and NO_2 , respectively, based on the full models: for nonaccidental mortality, HR = 1.08 (95% CI: 1.07, 1.09) and HR = 1.03 (95% CI: 1.02, 1.03); for mortality from respiratory diseases, HR = 1.13 (95% CI: 1.10, 1.17) and HR = 1.02 (95% CI: 1.01, 1.03); and for lung cancer mortality, HR = 1.26 (95% CI: 1.21, 1.30) and HR = 1.10 (95% CI: 1.09, 1.11). Only PM_{10} was associated with circulatory disease mortality (HR = 1.06; 95% CI: 1.04, 1.08 compared with HR = 1.00; 95% CI: 0.99, 1.01 for NO_2). For both pollutants, associations were strongest for lung cancer mortality (PM_{10} HR = 1.26; 95% CI: 1.21, 1.30; NO_2 HR = 1.10; 95% CI: 1.09, 1.11).

In the two-pollutant models, HRs for PM_{10} decreased for nonaccidental mortality and lung cancer mortality, and increased for

circulatory or respiratory disease mortality, still remaining statistically significant (Table 2). All HRs for NO_2 decreased after adjustment for PM_{10} , and the association with respiratory diseases was negative and no longer significant (Table 3).

To compare results with previous cohort studies, we calculated HRs for $\text{PM}_{2.5}$ assuming that the association with PM_{10} is driven by the $\text{PM}_{2.5}$ fraction (Beelen et al. 2014; Hoek et al. 2013). Based on a $\text{PM}_{2.5}/\text{PM}_{10}$ ratio of 0.66 (Cyrys et al. 2003; Eeftens et al. 2012) and PM_{10} HRs from the fully adjusted model, we estimated the following for each 10- $\mu\text{g}/\text{m}^3$ increase in $\text{PM}_{2.5}$: for nonaccidental mortality, HR = 1.13 (95% CI: 1.11, 1.14), for circulatory disease mortality HR = 1.09 (95% CI: 1.06, 1.12), for respiratory disease mortality HR = 1.18 (95% CI: 1.15, 1.27), and for lung cancer mortality HR = 1.41 (95% CI: 1.34, 1.49).

Associations between PM_{10} and nonaccidental, circulatory disease and lung

cancer mortality did not deviate significantly ($p < 0.01$) from linear (Figure 2). The association with respiratory disease mortality increased up to about 40 $\mu\text{g}/\text{m}^3$ and then unexpectedly decreased. Associations between NO_2 and mortality deviated significantly from linear only for circulatory disease (Figure 2).

In addition to our main analysis, we conducted a separate analysis of associations of PM_{10} and NO_2 with potential individual- and group-level confounders that were not available for the nationwide study sample, because the presence or absence of associations would help clarify the likelihood of confounding by these characteristics. After excluding data from adults < 30 years of age, information on lifestyle factors from 63,796 subjects was available, from 1,517 of the 3,985 four-digit postal code areas across the Netherlands. Age- and sex-adjusted mean PM_{10} and NO_2 for different categories of selected sociodemographic characteristics and lifestyle factors are presented in the




Figure 2. Estimated concentration–response curves (solid blue lines) and 95% CIs (dashed lines) for nonaccidental mortality, circulatory disease mortality, respiratory disease mortality, and lung cancer mortality for (A) NO_2 and (B) PM_{10} . In, natural logarithm. Model was adjusted for age, sex, marital status, region of origin, and household income.

Supplemental Material, Tables S1 and S2, respectively. PM_{10} concentrations differed statistically significantly between the various categories, although these differences were small ($< 0.2 \mu\text{g}/\text{m}^3$). After additional adjustment for marital status, region of origin, level of education, and neighborhood social status, these differences decreased to $0.07 \mu\text{g}/\text{m}^3$ for smoking and $0.10 \mu\text{g}/\text{m}^3$ for BMI, and 95% CIs overlapped (see Supplemental Material, Table S3). For NO_2 , these differences in age- and sex-adjusted mean concentrations were somewhat larger: $0.64 \mu\text{g}/\text{m}^3$ for current smokers compared with never smokers and 0.80 for $BMI > 30$ compared with $18.5 < BMI < 25$. After additional adjustment for marital status, region of origin, level of education, and neighborhood social status, these differences decreased to $0.46 \mu\text{g}/\text{m}^3$ both for smoking and for BMI, but remained statistically significant (see Supplemental Material, Table S3). When we evaluated differences in prevalence of lifestyle and sociodemographic characteristics across different categories (deciles) of PM_{10} and NO_2 exposure, the percentage of current smokers and participants with $BMI > 30$ are highest in the highest two deciles of PM_{10} and NO_2 exposure. However, the percentage of non-Dutch nationality, low education, and average neighborhood social status score follow a similar pattern, except for non-Dutch nationality in the highest decile of PM_{10} (see Supplemental Material, Table S4).

Additional adjustment for smoking-attributable mortality reduced the HR for PM_{10} by about 30% for non-accidental mortality (from 1.081 to 1.058) to 40% for the cause-specific mortalities (see Supplemental Material, Table S5). The HRs for NO_2 changed marginally for nonaccidental mortality (from 1.027 to 1.022), for deaths from diseases of the circulatory system (nonsignificant), and for lung cancer mortality (from 1.097 to 1.083). The NO_2 HR for respiratory mortality changed from 1.015 to a nonsignificant 1.003.

In Figure 3 we present the HRs for nonaccidental, circulatory, respiratory, and lung cancer mortality per $10 \mu\text{g}/\text{m}^3$ PM_{10} and NO_2 by sex, age, social economic status and degree of urbanization. We did not find consistent patterns of effect modification across the different outcomes, although, except for circulatory mortality, HRs tended to be closer to unity among those > 65 years of age compared with younger residents. For lung cancer, women were at higher risks for both, PM_{10} and NO_2 exposure.

Discussion

In this large Dutch nationwide population cohort of > 7 million adults we observed positive significant associations between estimated long-term exposure to air pollution

(PM_{10} and NO_2) at the home address and nonaccidental, circulatory disease, respiratory disease, and lung cancer mortality. We used large national demographic and geographical databases to assess these associations. Our results suggest that PM_{10} is more consistently associated with mortality than NO_2 .

The large size of the cohort allowed us to assess the small relative risks of ambient air pollution with more precision than in typical individual cohort studies. Interestingly, we found highly significant associations between PM_{10} and respiratory mortality. Associations between long-term exposure to fine particles and respiratory mortality have been inconsistent in previous individual cohort studies, partly due to a relatively small number of cases (Hoek et al. 2013). Our findings are in agreement with time-series studies based on registries that have shown consistent associations between day-to-day variation in air pollution and respiratory mortality. Consistent with the time-series studies, our estimates for respiratory mortality are larger than for nonaccidental mortality.

Another interesting finding is that although circulatory disease mortality was significantly associated with PM_{10} , hazard ratios were smaller than for nonaccidental mortality. Overall, in previous cohort studies, hazard ratios were larger for circulatory disease mortality, though with considerable variation between studies (Hoek et al. 2013). We speculate that better medication has reduced cardiovascular mortality, complicating assessment of associations with risk factors including air pollution.

Recently, several studies were published that were based on national demographic databases linked to air pollution data (Carey et al. 2013; Cesaroni et al. 2013; Chen et al. 2013; Crouse et al. 2012; Hales et al. 2012; Huss et al. 2010; Zeger et al. 2008). Three systematic reviews on the health effects of long-term exposure to air pollution (Brook et al. 2010; Chen et al. 2008; Hoek et al. 2013) give an extensive overview of the published literature on mortality and long-term exposure to air pollution. In Supplemental Material, Table S6, we summarize the results of the national registry studies and recent papers on mortality outcomes from a large European study on air pollution and health. For comparisons, we added our results into the table.

In our study we found particulate matter to be associated with all outcome measures that we have analyzed. Our relative risk estimate for PM_{10} on total mortality is higher than the relative risk estimate from a recent published study based on 19 European cohorts (Beelen et al. 2014). Only two other registry-based studies assessed the associations between mortality and PM_{10} (Carey et al.

2013; Hales et al. 2012). Both studies, like ours, reported higher HRs for respiratory mortality than for cardiovascular mortality (although the Hales study included lung-cancer mortality). A Dutch study by Beelen et al. (2008) also estimated higher HRs for respiratory mortality than for cardiovascular mortality (for $PM_{2.5}$), and Cesaroni et al. (2013) in Rome, Italy, reported higher HR for $PM_{2.5}$ for cardiovascular mortality than for respiratory mortality. For NO_2 we found statistically significant associations with all outcomes except for circulatory disease mortality, which is in line with earlier Dutch cohort study results (Beelen et al. 2008) in which a nonsignificant association for NO_2 was reported. Again, our estimate on total mortality is higher than the relative risk estimate from the recent published European study (Beelen et al. 2014). Our results for nonaccidental mortality are comparable with the results from the registry studies by Carey et al. (2013) and Cesaroni et al. (2013) and the Dutch cohort study by Beelen et al. (2008). Cause-specific results show more heterogeneity between the different studies presented in Supplemental Material, Table S6. In the analyses in which we adjusted associations with NO_2 for PM_{10} concentrations, the association between NO_2 and respiratory mortality disappeared while the PM_{10} effects remained. This suggests that mortality effects of particles are not explained exclusively by traffic, as was suggested in a recent European study as well (Beelen et al. 2014).

Our estimates for lung cancer mortality for PM_{10} and NO_2 are higher than those published recently in the national cohorts (Carey et al. 2013; Cesaroni et al. 2013; Hales et al. 2012; Huss et al. 2010), but comparable with the PM_{10} estimate for lung cancer incidence in a recently published study on the relation between long term exposure to air pollution and lung cancer incidence in 14 European cohorts (Raaschou-Nielsen et al. 2013). Adjustment for NO_2 levels reduced the estimate (HR = 1.093; 95% CI: 1.044, 1.144), whereas the estimate for NO_2 was HR = 1.080 (95% CI: 1.065, 1.096). This suggests that PM_{10} and NO_2 represent different characteristics of the air pollution mixture that act independently, which may be related to the source of the pollution variability. In the European study, no association with NO_2 was reported (Raaschou-Nielsen et al. 2013).

We found a tendency for lower HRs in the older age category (> 65 years). This is in line with the suggestion of effect modification by age in the Rome cohort (Cesaroni et al. 2013) and consistent with a previous study in Norway (Naess et al. 2007). In contrast with Cesaroni et al. (2013), we did not estimate stronger associations for men than women;




Figure 3. Adjusted hazard ratios (95% CIs) per 10-µg/m³ increase in PM₁₀ (A) and NO₂ (B), by population characteristics and cause of death. Model was adjusted for age, sex, marital status, region of origin, socioeconomic status, and household income, stratified as indicated on x-axis. Socioeconomic status categories are based on the quintiles of the social status rankings; urbanization is based on address density: (low: < 500 addresses/km²; moderate low: 500–1,000 addresses/km²; medium: 1,000–1,500 addresses/km²; moderate high: 1,500–2,500 addresses/km²; high: > 2,500 addresses/km²).

and for lung cancer, associations with both exposures were significantly stronger for women than men. We did not find clear evidence of modification by socioeconomic status, which is in line with the findings in the Rome cohort.

Although, in general, studies of long-term effects of air pollution on mortality qualitatively show similar results, differences in the quantitative outcomes remain.

Our results contribute to the evidence linking long-term ambient air pollution exposure to increased nonaccidental and cause-specific mortality. Our study has several strengths. The study size is very large and includes all Dutch citizens of ≥ 30 years of age in 2004, living at least for 5 years at the 2004 address, which improves the long-term exposure classification. We used address level for all cohort members, based on the annual mean NO_2 and PM_{10} concentrations at a $100 \text{ m} \times 100 \text{ m}$ grid. Further, because of the relatively low correlation between the two air pollutant components ($R = 0.58$), we were able to disentangle the relative importance of the two components when adjusting for each other. We had individual information about important predictors of mortality at both the individual and ecological levels.

Apart from the strengths, there are also limitations of our study. Exposures were estimated by a LUR model for the year 2001 and assigned to the follow-up period 2004–2011. Although the exposure assignment precedes the follow-up period, we are not sure that the 2001 annual average adequately represents a longer exposure window, which is relevant for long-term exposure. However, there is evidence from the literature that spatial distribution of air pollution is stable over 10-year periods (Cesaroni et al. 2012; Eeftens et al. 2011; Gulliver et al. 2011). Still, people might have moved since 2004 to unknown addresses and therefore changed their exposure.

Because of the limited availability of individual lifestyle factors, we adjusted for individual and area-level SES, as in the American Cancer Society study (ACS) (Krewski et al. 2009) and the European Study of Cohorts for Air Pollution Effects (ESCAPE) (Beelen et al. 2014). The two confounders represent different “contextual” environments. Overadjustment is a possibility, but unlikely, because the correlation between the two covariates was low (-0.11).

No information on individual risk factors such as smoking, diet, alcohol use, and obesity was available. We evaluated the possibility that uncontrolled confounding from lifestyle factors may have biased our results using data from $> 60,000$ 30- to 65-year-old participants of health surveys conducted in 2003–2005 by 11 Community Health Services. We do not have appropriate

references to substantiate the representativeness of this subgroup in comparison to our national cohort, but because the regions were spread over the Netherlands and the goal of the surveys is to select random samples, we think that exposure ranges are well covered in the subgroup and comparable with the exposure range in the national cohort.

After adjustment for age, sex, marital status, region of origin, level of education, and neighborhood social status score, mean PM_{10} and NO_2 exposures were only $0.1 \mu\text{g}/\text{m}^3$ (PM_{10}) and $0.5 \mu\text{g}/\text{m}^3$ (NO_2) higher among current smokers compared with never smokers, and among participants with $\text{BMI} > 30$ compared to $18.5 < \text{BMI} < 25$. When looking at the prevalence of lifestyle and sociodemographic characteristics across different categories (deciles), we found that these differences were likely driven by a higher percentage of current smokers and participants with high (> 30) BMI in the highest two deciles of PM_{10} and NO_2 exposure.

As we observe associations for the whole exposure distribution, we think it is unlikely that uncontrolled confounding from smoking or BMI has substantially biased our results (Figure 2; see also Supplemental Material, Tables S1–S4.)

In a second sensitivity analysis, we used regional age-standardized smoking attributable mortality fractions, 22–30% for men and 7–14% for women, in 40 NUTS-3 (nomenclature of territorial units for statistics) regions (Janssen and Spriensma 2012). The average population in these regions is about 400,000. We found that additional adjustment in the analyses with an area-level proxy for smoking reduces the HRs but did not materially affect the conclusions of the study (see Supplemental Material, Table S5). Several cohort studies have been published with missing data on, presumably, important individual confounders (see Supplemental Material, Table S6), but most of them showed that relative risk estimates did not materially change when some additional adjustment was made by using proxies for individual risk factors on an aggregated level (Cesaroni et al. 2013; Chen et al. 2013; Villeneuve et al. 2011; Zeger et al. 2008). In general, this suggests evidence that the lack of individual data on smoking and BMI did not bias the results in such a way that conclusions drawn on data with missing information for some individual potential confounders are materially wrong. Krewski et al. (2005a, 2005b) showed that adjusting for self-reported smoking had little effects on the relative risk estimates in two different U.S. cohorts. In the National English Cohort (Carey et al. 2013) the effect estimates were robust to adjustment for individual smoking and BMI (collected from individual patient

cohorts). This was also acknowledged by Hoek et al. (2013), who concluded that effect estimates from the three large population cohorts without individual smoking data (Cesaroni et al. 2013; Crouse et al. 2012; Zeger et al. 2008) were not higher than those from the individual cohort studies.

Conclusions

Long-term exposure to particulate air pollution (PM_{10}) and NO_2 was associated with nonaccidental mortality, mortality from respiratory diseases, and lung cancer mortality in our study population of 7 million adults. Furthermore, PM_{10} was associated with cardiovascular mortality. Associations with PM_{10} were robust to adjustment for NO_2 , and associations of NO_2 with nonaccidental and lung cancer mortality remained after adjustment for PM_{10} .

REFERENCES

- Abbey DE, Nishino N, McDonnell WF, Burchette RJ, Knutsen SF, Lawrence Beeson W, et al. 1999. Long-term inhalable particles and other air pollutants related to mortality in nonsmokers. *Am J Respir Crit Care Med* 159:373–382.
- Arts CH, Hoogteijling EMJ. 2002. Het Sociaal Statistisch Bestand 1998 en 1999 [in Dutch]. Sociaal-economische Maandstatistiek 12:13–21.
- Beelen R, Hoek G, van den Brandt PA, Goldbohm RA, Fischer P, Schouten LJ, et al. 2008. Long-term effects of traffic-related air pollution on mortality in a Dutch cohort (NLCS-AIR study). *Environ Health Perspect* 116:196–202; doi:10.1289/ehp.10767.
- Beelen R, Raaschou-Nielsen O, Stafoggia M, Andersen ZJ, Weinmayr G, Hoffmann B, et al. 2014. Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the multicentre ESCAPE project. *Lancet* 383:785–795.
- Brook RD, Rajagopalan S, Pope CA III, Brook JR, Bhatnagar A, Diez-Roux AV, et al. 2010. Particulate matter air pollution and cardiovascular disease: an update to the scientific statement from the American Heart Association. *Circulation* 121:2331–2378.
- Brunekreef B, Beelen R, Hoek G, Schouten L, Bausch-Goldbohm S, Fischer P, et al. 2009. Effects of long-term exposure to traffic-related air pollution on respiratory and cardiovascular mortality in the Netherlands: the NLCS-AIR study. *Res Rep Health Eff Inst* 139:5–71.
- Carey IM, Atkinson RW, Kent AJ, van Staa T, Cook DG, Anderson HR. 2013. Mortality associations with long-term exposure to outdoor air pollution in a national English cohort. *Am J Respir Crit Care Med* 187:1226–1233.
- Cesaroni G, Badaloni C, Gariazzo C, Stafoggia M, Sozzi R, Davoli M, et al. 2013. Long-term exposure to urban air pollution and mortality in a cohort of more than a million adults in Rome. *Environ Health Perspect* 121:324–331; doi:10.1289/ehp.1205862.
- Cesaroni G, Porta D, Badaloni C, Stafoggia M, Eeftens M, Meliefste K, et al. 2012. Nitrogen dioxide levels estimated from land use regression models several years apart and association with mortality in a large cohort study. *Environ Health* 11:48; doi:10.1186/1476-069X-11-48.
- Chen H, Goldberg MS, Burnett RT, Jerrett M, Wheeler AJ, Villeneuve PJ. 2013. Long-term

- exposure to traffic-related air pollution and cardiovascular mortality. *Epidemiology* 24:35–43.
- Chen H, Goldberg MS, Villeneuve PJ. 2008. A systematic review of the relation between long-term exposure to ambient air pollution and chronic diseases. *Rev Environ Health* 23:243–297.
- Crouse DL, Peters PA, van Donkelaar A, Goldberg MS, Villeneuve PJ, Brion O, et al. 2012. Risk of nonaccidental and cardiovascular mortality in relation to long-term exposure to low concentrations of fine particulate matter: a Canadian national-level cohort study. *Environ Health Perspect* 120:708–714; doi:10.1289/ehp.1104049.
- Cyrys J, Heinrich J, Hoek G, Meliefste K, Lewné M, Gehring U, et al. 2003. Comparison between different traffic-related particle indicators: elemental carbon (EC), PM_{2.5} mass, and absorbance. *J Expo Anal Environ Epidemiol* 13:134–143.
- de Bruin A, Kardaun JW, Gast A, de Bruin E, van Sijl M, Verweij G. 2004. Record linkage of hospital discharge register with population register: experiences at Statistics Netherlands. *Stat J UN Econ Comm Eur* 21:23–32.
- Dockery DW, Pope CA III, Xu X, Spengler JD, Ware JH, Fay ME, et al. 1993. An association between air pollution and mortality in six U.S. cities. *N Engl J Med* 329:1753–1759.
- Eeftens M, Beelen R, de Hoogh K, Bellander T, Cesaroni G, Cirach M, et al. 2012. Development of Land Use Regression models for PM_{2.5}, PM_{2.5} absorbance, PM₁₀ and PM_{coarse} in 20 European study areas; results of the ESCAPE project. *Environ Sci Technol* 46:11195–11205.
- Eeftens M, Beelen R, Fischer P, Brunekreef B, Meliefste K, Hoek G. 2011. Stability of measured and modelled spatial contrasts in NO₂ over time. *Occup Environ Med* 68:765–770.
- GGD GHOR Nederland. 2014. GGD GHOR Nederland Homepage. Available: <http://www.ggdghor.nl/english/> [accessed 16 February 2015].
- Gulliver J, Morris C, Lee K, Vienneau D, Briggs D, Hansell A. 2011. Land use regression modeling to estimate historic (1962–1991) concentrations of black smoke and sulfur dioxide for Great Britain. *Environ Sci Technol* 45:3526–3532.
- Hales S, Blakely T, Woodward A. 2012. Air pollution and mortality in New Zealand: cohort study. *J Epidemiol Community Health* 66:468–473.
- Harteloh P, de Bruin K, Kardaun J. 2010. The reliability of cause-of-death coding in the Netherlands. *Eur J Epidemiol* 25:531–538.
- Hoek G, Krishnan RM, Beelen R, Peters A, Ostro B, Brunekreef B, et al. 2013. Long-term air pollution exposure and cardio-respiratory mortality: a review. *Environ Health* 12:43; doi:10.1186/1476-069X-12-43.
- Huss A, Spoerri A, Egger M, Rössli M, Swiss National Cohort Study Group. 2010. Aircraft noise, air pollution, and mortality from myocardial infarction. *Epidemiology* 21:829–836.
- Janssen F, Spiensma AS. 2012. The contribution of smoking to regional mortality differences in the Netherlands. *Demogr Res* 27:233–260.
- Knol FA. 1998. Van hoog naar laag, van laag naar hoog: de sociaal-ruimtelijke ontwikkeling van wijken tussen 1971-1995 [in Dutch]. The Hague, the Netherlands: Sociaal en Cultureel Planbureau.
- Krewski D, Burnett RT, Goldberg M, Hoover K, Siemiatycki J, Abrahamowicz M, et al. 2005a. Reanalysis of the Harvard Six Cities Study, part 1: validation and replication. *Inhal Toxicol* 17:335–342.
- Krewski D, Burnett RT, Goldberg M, Hoover K, Siemiatycki J, Abrahamowicz M, et al. 2005b. Reanalysis of the Harvard Six Cities Study, part 2: sensitivity analysis. *Inhal Toxicol* 17:343–353.
- Krewski D, Jerrett M, Burnett RT, Ma R, Hughes E, Shi Y, et al. 2009. Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality. *Res Rep Health Eff Inst* 140:5–114.
- Naess Ø, Nafstad P, Aamodt G, Clausen B, Rosland P. 2007. Relation between concentration of air pollution and cause-specific mortality: four-year exposures to nitrogen dioxide and particulate matter pollutants in 470 neighborhoods in Oslo, Norway. *Am J Epidemiol* 165:435–443.
- Pope CA III, Thun MJ, Namboodiri MM, Dockery DW, Evans JS, Speizer FE, et al. 1995. Particulate air pollution as a predictor of mortality in a prospective study of U.S. adults. *Am J Respir Crit Care Med* 151(3 pt 1):669–674.
- Prins CJM. 2000. Dutch population statistics based on population register data. *Maandstat Bevolking* 48:9–15.
- R Core Team. 2014. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing. Available: <http://www.R-project.org> [accessed 16 February 2015].
- Raaschou-Nielsen O, Andersen ZJ, Beelen R, Samoli E, Stafoggia M, Weinmayr G, et al. 2013. Air pollution and lung cancer incidence in 17 European cohorts: prospective analyses from the European Study of Cohorts for Air Pollution Effects (ESCAPE). *Lancet Oncol* 14:813–822.
- van Elzakker BG, Buijsman E. 2014. The Dutch National Air Quality Monitoring Network: Monitoring Program in 1999. Available: http://www.rivm.nl/en/Documents_and_publications/Scientific/Reports/1999/maart/The_Dutch_National_Air_Quality_Monitoring_Network_monitoring_program_in_1999?sp=cml2bXE9ZmFsc2U7c2VhcmNoYmFzZT01OTAxMDtyaXZtcT1mYWxzZTs=&page=5902 [accessed 16 February 2015].
- Vienneau D, de Hoogh K, Beelen R, Fischer P, Hoek G, Briggs D. 2010. Comparison of land-use regression models between Great Britain and the Netherlands. *Atmos Environ* 44:688–696.
- Villeneuve PJ, Goldberg MS, Burnett RT, van Donkelaar A, Chen H, Martin RV. 2011. Associations between cigarette smoking, obesity, sociodemographic characteristics and remote-sensing-derived estimates of ambient PM_{2.5}: results from a Canadian population-based survey. *Occup Environ Med* 68:920–927.
- Yap C, Beverland IJ, Heal MR, Cohen GR, Robertson C, Henderson DE, et al. 2012. Association between long-term exposure to air pollution and specific causes of mortality in Scotland. *Occup Environ Med* 69:916–924.
- Zeger SL, Dominici F, McDermott A, Samet JM. 2008. Mortality in the Medicare population and chronic exposure to fine particulate air pollution in urban centers (2000–2005). *Environ Health Perspect* 116:1614–1619; doi:10.1289/ehp.11449.